




Einstein to Google: New Approaches to Dark Energy



Einstein Equations


$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi G T_{\mu\nu}$$

GEOMETRY

Expansion Rate


ENERGY

Potential Energy, proportional to density




Observed acceleration points to dark energy that remains roughly constant


Dark Energy Task Force (2006)



Supernova Brightness




Baryon Acoustic Oscillations




Gravitational Lensing





Galaxy Cluster
Abundance






Crack #1: Probes are Correlated







Crack #2: Maybe Einstein was wrong

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R \neq 8\pi G T_{\mu\nu}$$

Einstein was smart, and coming up with a better theory of gravity is hard

Crack #2: Maybe Einstein was wrong

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R \neq 8\pi G T_{\mu\nu}$$


Einstein was smart, and coming up with a better theory of gravity is hard

Modified Gravity vs. Dark Energy + GR


- New degrees of freedom
- MG models can postdict any expansion history
- With fixed expansion history, structure grows differently in the two classes of models

Population of the U.S.


- Total population grew from 4M->310M
- Pattern of over- and underregions grew in space and time
- If we understood all the forces –
 sociological, economic, racial,
 and political -- acting on the
 people ... we would be able to
 explain these patterns and this
 evolution




Schematically: Fix Expansion History and Measure Growth of Structure




Snowmass: Huterer et al. 2013





BAO and **SN** measure expansion history, but **Clusters** and **Lensing** sensitive to both expansion history and growth. All are correlated. Many different options for data vector.







Can compute for Dark Energy model with (w,w') but many different MG models exist. Also need to include small scale modeling. Many different options for cosmological and nuisance parameters.










Covariances depend on theory; require many simulations. Many different approaches even for fixed data vector, parameters






Will need to include priors from CMB and other experiments. Many different choices







Cosmological Survey Inference System

(CosmoSIS) Funded by: DOE Computational HEP & NSF Physics Frontier Center











Marc Paterno

Jim Kowalkowski, Scott Dodelson

Saba Sehrish







Sarah Bridle

Joe Zuntz









Doug Rudd

Elise Jennings Alessandro Manzotti

Break-out workshop (25 people) May, 2014; Sessions at DES and upcoming LSST collaboration meetings

CosmoSIS
Standard
Library:
camb, wmap,
planck, cfht,
bicep, ...

DES:

C_l, D(z), MG models, joint likelihoods, nuisance parameters, covariances, halo model, ...

LSST

Delivered in ups packages and repos

CosmoSIS
Core:
Interfaces,
Build systems,
run time

Python, scipy, gcc, gsl, fftw, ...

Think of us as your younger sibling



Some New Features



- Cosmosis-standard-library has contributions from other probes, experiments, etc.
- Can upload modules in multiple languages
- Credit for uploaded modules in "terms of use"
- Parallelism: natural to deploy and run on an HPC cluster

Einstein-> Google

- Modified Gravity may explain acceleration
- Four traditional DE probes highly correlated
- Many ways of answering key physics questions: What is the nature of dark energy?
 Is it dark energy or modified gravity?
- CosmoSIS framework that will help thousands of scientists work on these questions together